

This is Google's cache of http://www.preparedfoods.com/archives/2002/2002_3/0302formulation.htm.

Google's cache is the snapshot that we took of the page as we crawled the web.

The page may have changed since that time. Click here for the [current page](#) without highlighting.

To link to or bookmark this page, use the following url:

http://www.google.com/search?q=cache:jOmgAgB7kNYC:www.preparedfoods.com/archives/2002/2002_3/0302formulation.htm+soybean+oil+ph

Google is not affiliated with the authors of this page nor responsible for its content.

These search terms have been highlighted: **soybean oil phospholipid lecithin**

**PREPARED
FOODS.com**

[Home page](#)

Browse

[by introduction](#)

[by subject](#)

[by issue](#)

2002 20th Annual New Products Conference
October 13-16, 2002
Fairmont Scottsdale Princess Resort
Scottsdale, Arizona

» [March 2002 Issue](#) Formulation & Ingredient Challenges

Boosting Bread's Value

Emulsifiers are key to baked goods' texture and shelf life. **Lecithin and soybread, in particular, are profiled. Ingredients from fat, oil and emulsifier suppliers are showcased.**

Laura A. Brandt
Technical Field Editor

Fats, oils and emulsifiers play an important role in baked goods. For example, in soy-based breads, soy flour absorbs water to form a gel, making the loaves quite dense. Emulsifiers help open up the bread texture for less-dense loaves, with greater volume. In addition, emulsifiers improve dough handling and help extend bread's shelf life.

Emulsifiers

By using a surfactant, such as an emulsifier, one can create an emulsion of two or more incompatible substances. A typical emulsion is a dispersion of small droplets of one immiscible liquid within another. Emulsifiers keep the droplets from coalescing. These ingredients not only offer emulsification, but also starch complexing, protein strengthening and aeration in baked goods.

Emulsifiers are used in bread to extend shelf life by slowing down the staling process. Bread tastes and feels stale when it starts to taste like cardboard, and feels dry and hard. In the laboratory, staleness can be measured by the crumb's resistance to compression, with an instrument such as the Instron Universal Testing Machine.




Emulsifiers used in breads include mono- and di-glycerides, various forms of lecithin, diacetyl tartrate ester of monoglyceride (DATEM), and sodium stearoyl lactylate (SSL). Most bakeries use a blend of these functional ingredients.

According to research studies, staling is not related to moisture loss from bread. While days old bread gives a drier mouthfeel, it has the same moisture level as fresh bread.

Rather than producing an initially softer crumb, as previously believed, emulsifiers interfere with the recrystallization (or retrogradation) of amylose, which retards the firming rate. The fatty acid-end of emulsifiers forms a complex with gelatinized starch to slow down starch recrystallization and the staling process in bread. (See chart on page 42.)

Emulsifiers used in breads include mono- and di-glycerides, various forms of lecithin, diacetyl tartrate ester of monoglyceride (DATEM), and sodium stearoyl lactylate (SSL). Emulsifiers have different functions in bread, e.g. lecithin behaves as a wetting agent, while DATEM and SSL act as dough strengtheners. Most bakeries use a blend of these functional ingredients.



The rate of staling in wheat bread can be quantified by the amount of starch molecule (amylopectin) recrystallization, as measured by DSC (differential scanning calorimetry). Adding soy decreases recrystallization and, thus, retards staling, as compared to a control without soy.

Lecithin

Lecithin, a primary emulsifier in egg yolks, also can be derived from rice and other grains. Soy-based lecithin, which contains phospholipids, is a by-product from the refining of crude soybean oil. The four main

components of soy lecithin include phosphatidylcholine, phosphatidylethanolamine, phosphatidylinositol, and phosphatidic acid. Soy-based lecithin has a more healthful image than some other emulsifiers.

With lecithin's hydrophobic and hydrophilic portions, it can simultaneously interact with both oil and water, making it an effective emulsifier. In addition, lecithin aids in machinability and shortening dispersion of baked goods. Various types of lecithin are available to bakers including fluid, deoiled (hydrophilic), chemically-modified (acetylated/hydroxylated), enzyme-modified (EM) and powdered versions of these.

Considerations for selecting lecithin are specific benefits, labeling requirements and handling procedures. Fluid lecithin is a viscous, syrupy, amber-colored liquid. Bakers prefer powdered lecithin. It is easy to incorporate with dry ingredients and is convenient to add to dry mixes.

Powdered, deoiled lecithin, commonly used in breads, has less flavor than fluid lecithin, is higher in phospholipid content, and is water dispersible. Standard fluid lecithin contains about 30% soybean oil, with deoiled lecithin containing 2% or less, says Karen Allen-Seabolt, associate scientist with Central Soya Co., Inc., Lecithin Group, Fort Wayne, Ind. Usage level is about 0.5 to 1.0% on a flour weight basis.

"We have different forms of lecithin to help increase volume and dough strengthening, and to extend shelf life," says Seabolt. "Deoiled, non-modified lecithin helps improve crumb structure and volume, but it doesn't help as much as enzyme-modified lecithin with shelf life for yeast-raised products."

Deoiled lecithin is more concentrated in phospholipids—the active, functional component of lecithin—than fluid forms, according to Bruce Seabee, Ph.D., manager of Emulsifier and Texturant Technology, ADM Research, Decatur, Ill. In most formulations, deoiled lecithin can replace standard fluid at two-thirds usage rate, he adds. EM lecithin performs similarly to monoglycerides because it will form a complex with starch more readily.

"In enzyme-modified lecithin, the middle-position fatty acid is removed from the phospholipid with a phospholipase enzyme," says Seabolt. "It is replaced with a hydroxyl group, making the molecule more polar, which increases its ability to complex with starch. EM lecithin is used mainly for yeast-raised baked goods to extend shelf life, but it also improves volume with its ability to act as a dough strengthening agent."

EM lecithin can replace mono-glycerides. Its use in bread is becoming more popular with bakeries today, adds Seabolt.

Alvarado Street Bakery uses several soy-based ingredients in its Sprouted Soy Crunch Bread, including: Organically-grown sprouted soybeans, toasted soy nuts, organically-grown whole soy flour, soy-based lecithin, as well as unsulphured molasses and organically-grown sprouted whole wheat. The bread is frozen, then shipped to mainstream supermarkets and health food stores coast to coast.



Website Resources

www.preparedfoods.com/archives/1997/1854.htm – PF article overview on science of emulsifiers and processing
www.scisoc.org/aacc/meeting/2001/abstracts/a01ma350.htm
– AACC abstract on consumer acceptability of soy bread
www.eurekalert.org/pub_releases/2001-12/osum-oss120601.php – Press release on heart-healthy soy bread
www.frenchmeadow.com/ – French Meadow Bakery, Minneapolis, Minn.
www.alvaradostreetbakery.com/ – Alvarado Street Bakery, Rohnert Park, Calif.
www.preparedfoods.com/archives/2001/2001_4/0401soyfoodssoar.htm – PF article on soyfood
www.scisoc.org/aacc/meeting/2001/abstracts/a01ma375.htm
– AACC abstract on soy's effect on bread properties during storage

OSU Researchers Formulate Tasty Soy Bread

Rather than relying on a nutritional bar or a beverage, researchers at Ohio State University (OSU) are hoping that consumers will consider an everyday food to get soy into their diets—bread.

One large slice or two slices (50g) of the bread contains 6.25g of soy protein per serving, which meets the soy protein health claim criteria.

"I didn't start looking for health claims—I became interested in adding soy to bread formulations and comparing wheat bread and soy breads because staling seemed to be retarded with the addition of soy flour to wheat flour," says Yael Vodovotz, assistant professor of food science at Ohio State University and co-author of the book, *Bread Staling*. (CRC Press, 2000).




Vodovotz found adding soy to bread decreases amylopectin recrystallization significantly, increasing the soy bread's shelf life.

The bread contains flour made from hard red wheat and two soy-based ingredients—soy flour and soymilk powder. While loaf volume decreases with an increase in soy flour, soymilk helps increase the loaf volume and has a positive flavor impact, says Vodovotz. Adding mono- and di-glycerides—as well as some fat—helps increase the loaf volume by lubricating the dough system and increasing the flexibility of the intercellular structure. A dough strengthener, L-cysteine, boosts internal structure.

A consumer acceptability study by Cory Ballard, an OSU undergraduate and senior food technologist for Chef-Con, Delaware, Ohio, found that 70% of those who participated in in-store sensory tests preferred the flavor of soy bread to that of red wheat bread. "Kids really liked the soy bread because it's sweet," says Ballard.

In another study A.S. Colakoglu, an OSU student, investigated "The effect of soy on the thermal and mechanical properties of bread during storage." Differential Scanning Calorimeter (DSC) studies showed that the addition of soy flour (20%-40% substitution) to wheat flour decreased the amount of "freezable" water and amylopectin recrystallization, which can be interpreted to say that adding soy flour retards staling.



can be interpreted to say that adding soy flour retards staling.

OSU is licensing the formula to bakeries for sale in mainstream supermarkets. Researchers at the OSU medical school want to test the bread's cancer-control potential.

➤ [Back to top](#)

This is **G o o g l e**'s cache of http://www.goldenflax.com/home_pages/omega3.htm.

G o o g l e's cache is the snapshot that we took of the page as we crawled the web.

The page may have changed since that time. Click here for the current page without highlighting.

To link to or bookmark this page, use the following url:

http://www.google.com/search?q=cache:eSRp51A4O00C:www.goldenflax.com/home_pages/omega3.htm+omega-3+sunflower&hl=en&ie=UTF-8

Google is not affiliated with the authors of this page nor responsible for its content.

These search terms have been highlighted: **omega 3 sunflower**



health benefits

[email](#) [home](#) [to order](#)

nutrition
information

flax facts
fiber • lignans • omega3

preparing flax

flaxseed vs.
flax oil

whole vs.
milled flaxseed

where can I
purchase flax?

research &
information links

frequently

golden flaxseed
omega-3

Society in general has become overweight, and we immediately tend to jump on the bandwagon of condemning the consumption of foods categorized in the "fats" food group. Even the term "fats and oils" will sometimes send shivers of repulsion into the minds of those who maintain a rigid and uncompromising diet pattern, swearing off all foods from this necessary food group. "Fats" has somehow become the shunned four-letter word of dietary watchdogs. But in our quick dismissal of trimming fat from our diets, we have also tended to forget some of the necessary functions of what is known as "essential fats". The terminology is certainly apt; as a certain amount of fat in the diet is absolutely essential for our health. Learning about "good fat" and "bad fat" is crucial to assist in making healthy choices. Cutting out all fat in your diet is not a prudent or healthy choice.

From Cave Dwellers to Cond -Own rs

Civilization has certainly changed our lifestyles. Yet, the basic bodily needs have not changed since the beginning. Balance in life again is key. Our health

asked questions

photo comparison

about north
american nutrition

▶order

golden flax seed

web designed by
Allegromkt .com

beginning. Balance in life again is key. Our health relies upon the basic needs of fresh air, clean water, sunshine, exercise and a common sense balance of proteins, carbohydrates and fats.

Our diet has changed significantly since our cave-dwelling ancestors inhabited the earth. Certainly the "good old days" are not always as good as one tends to conceive. Yet, studies of the hunter-gatherer lifestyle have shown innate understandings of the body's need for a balanced diet. This is where we gain even greater persuasion to understand the importance of essential fatty acids (EFA).

Due to lifestyle and dietary habits with fast-food and convenience foods, the general public has consumed a diet rich in fats. A common reaction then is to eliminate all or the majority of fat in the diet which is equally detrimental. The problem lies in the type of fat we ingest –too much saturated fat - known as the "bad fats" - the type that clogs arteries and raises cholesterol levels.

The diet of our ancestors included a good balance of essential fatty acids (EFA). Essential fatty acids (EFA) are grouped into two families, the omega-6 EFAs and the **omega-3** EFAs. The omega groups are considered the "good fats" or known as the polyunsaturated fats. The omega-6 fatty acids are found in corn, safflower, **sunflower**, canola and soybean oils. **Omega-3** EFAs are found in flaxseeds and flaxseed oil along with fish such as salmon, herring, trout, sardines and albacore tuna. Ground flaxseed and flaxseed oil provide a natural and concentrated level of **omega-3** EFAs, without concern for chemical contamination that may be a risk with fish consumption.

Finding Balance – The Alpha and the Omega

Understanding the difference between the omega-6 and the **omega-3** fatty acid is very important to know. It is more than just the difference in their numbers that should concern you.

Omega-6 fatty acids (found in vegetable oils with high proportions of linolenic acid) are best used by the body in a range of anywhere from a 4:1 to a 1:1 proportion with the **omega-3** fatty acids. We need both **omega-3** and omega-6 fatty acids. Yet, an excess of omega-6 fatty acids can have dire consequences. Many

scientists believe that a major reason for the high incidence of heart disease, hypertension, diabetes, obesity, and some forms of cancer is the extreme imbalance between our intake of omega-6 and **omega-3** fatty acids.

The parent compound in the **omega-3** fatty acid is called alpha-linolenic acid (ALA). It is this compound that serves as the "computer" or brain for the **omega-3** fatty acid in determining how it will best maximize the body's functioning.

Our ancestors evolved on a diet with a 1:1 ratio of omega-6 to **omega-3** fatty acids. Dietary changes over the last few centuries have changed this ratio anywhere from 20:1 to 25:1. This is clearly an equation for trouble, and today's chronic health problems obviously exemplify this concern.

One of the primary reasons we ingest too much of the omega-6 fatty acid groups in our diet is the mass use of vegetable oils. This practice is so far-reaching that practically every fried food and snack food available has been cooked in soybean, corn, **sunflower** or canola oil. These oils are usually processed by hydrogenation. This changes their molecular structure so they are basically good for frying foods at a high temperature and providing a lengthy shelf-life in the grocery store. Unfortunately, these molecular properties in the omega-6 fatty acids promote inflammation, blood clotting and tumor growth.



The **omega-3** fatty acids act entirely opposite. But, when the omega-6 fatty acids are disproportionately higher, the **omega-3** fatty acids cannot compete with the omega-6 activity. When in balance, they work in concert, making sure for every action there is a reaction, helping to maintain stability in the body.

When the omega-6 and **omega-3** fatty acids maintain a healthy balance; they effectively become clearinghouses or message centers to the rest of the body to:

- Alert the immune system to go into action
- Signal the blood vessels to either widen or narrow
- Tell blood platelets to clot or not by sticking together or separating
- Regulate inflammation

- Formulate neural networks for brain activity in learning, memory processes and mood regulation

Trouble is brewed when one fatty acid overpowers another. Clearly, the data shows we need to seriously increase **omega-3** fatty acids in our diets. **Omega-3** fatty acid deficiencies are increasingly prevalent with young children. A Purdue University study showed that children low in **omega-3** essential fatty acids are significantly more likely to be hyperactive, have learning disorders and to display behavioral problems.

In the general public, studies have linked **omega-3** deficiencies to chronic health problems of diabetes, cancer, arthritis, inflammatory diseases, depression, heart disease, hypertension, memory problems, weight gain and some allergies and skin conditions.

Researchers believe 60% of Americans are deficient in **omega-3** fatty acids and approximately 20% of those have so little that test methods would not be able to detect even a trace in their blood.

Changing the Scenario

We imagine you are now convinced of the need for adding **omega-3** fatty acids to your diet. Yet, probably wondering how to go about doing it and also wondering how much you need to add. As noted prior, the two major sources of **omega-3** fatty acids are fish such as salmon, trout, and albacore tuna and flaxseed. Due to high risk of chemical contamination in fish products, we recommend flaxseed and flaxseed products to boost your **omega-3** fatty acid consumption. Flaxseed is loaded with alpha-linolenic acid. Remember this is the "brain" of the **omega-3** fatty acid molecule and assists in maximizing the benefits of nutritious foods. Most foods have far less **omega-3** properties than what is found in flaxseed. In fact, it would take 25 cups of peanut butter to get the alpha-linolenic acid found in just _ cup of ground flaxseed. Imagine those calories, not to mention the stares you might get when going through the grocery line with a cart full of peanut butter! To gain further information about adding flaxseed to your diet you can link to the [recipes](#) and [preparing flax sections](#). Helpful facts are also provided in the [nutrition information](#), with accompanying dietary guidelines.



To order Omega3 Golden Flax online - click here.

NOTE: Information presented here does not replace seeking advice from your physician.

[email](#) - [home](#) - [health benefits](#) - [nutrition information](#) - [flax facts](#) -
[preparing flax](#) - [flaxseed vs. flax oil](#) - [whole vs. milled flaxseed](#) -
[where can i purchase flax?](#) - [research & information links](#) -
[frequently asked questions](#) - [photo comparison](#) - [to order](#)

This is Google's cache of http://www.ivillage.co.uk/health/comp/herb/qas/0,12712,181042_182146,00.html.

Google's cache is the snapshot that we took of the page as we crawled the web.

The page may have changed since that time. Click here for the [current page](#) without highlighting.

To link to or bookmark this page, use the following url:

http://www.google.com/search?q=cache:YUv1CW8Y3SOC:www.ivillage.co.uk/health/comp/herb/qas/0,12712,181042_182146,00.html+evening+

Google is not affiliated with the authors of this page nor responsible for its content.

These search terms have been highlighted: **evening primrose oil omega**

with **one** Personal Loan from
Marks & Spencer Financial Services

consolidate
your debt

iVillage.co.uk

you are here ► [iVillage.co.uk](#) ► [health & beauty](#) ► [complementary health](#) ► [herbs & supplements](#)

home

join free • email • message boards • horoscopes • quizzes • search

iVillage UK

health & beauty

▼ topics

choose...

choose... iVillage Home
Computers Diet&Fitness
Food Games
Health&Beauty Horoscopes
Money Motoring
News&Politics
Pregnancy&Baby
Relationships Shopping
Work&Career

▼ categories

Women's health
General health
Complementary health
Healthy living
Ages and stages
Beauty
Disorders A - Z
Auctions

▼ support and advice

- Let's talk about health
- Breast cancer support
- Coping with depression
- Giving up smoking
- Beauty talk

▼ tools and quizzes

- Stain solver
- Men's health quiz

Evening primrose oil: what's all the fuss about?

by Alyson Greenhalgh and Dr. Sara Kirk

Q

What does evening primrose oil do?

A

Evening primrose oil is a nutritional supplement and a member of the essential fatty acid (EFA) group. There are two members of the group, the **omega-3 and omega-6 fatty acids**.

Evening primrose oil is rich in **omega-6 fatty acids** and one of these, **gamma linolenic acid (GLA)**, has certain medicinal properties that may act as an anti-inflammatory for conditions such as arthritis, eczema, high blood pressure or hypertension. Evening primrose oil is also believed to help alleviate cramps associated with pre-menstrual symptoms, though this has yet to be scientifically proven. Eating fatty fish such as tuna, mackerel, salmon and herring, and using safflower, sunflower and rapeseed oils in cooking is the first

You and your health

[Stop Smoking Challenge](#)
[Let's talk about health](#)
[Use the Health Calculator](#)
[Try our beauty tips](#)
[Learn about general health](#)
[Treat yourself with herbs](#)
[Stop Smoking SMS Support](#)
[Stop Smoking Quiz](#)
[Stop Smoking Message Board](#)
[Think about that diet!](#)

- Are you sun smart?
- Banish the blemish
- Health calculator
- Stop smoking quiz
- Allergies quiz
- Cold or flu quiz
- Migraine quiz
- Keyboard yoga
- Are you burning out?
- Free IQ Report

▼ experts corner

- Hair Expert NEW
- General Practitioner
- Nutritionist
- Complementary Health Expert

▼ services

- Online Dieting
- Stop Smoking Tips
- Send an e-Card
- Phone Book
- Competitions

▼ Free newsletters

Astrology Baby and Toddler
Beauty Diet and Fitness Food
Health Pregnancy
Relationships Work and
Career iVillage UK

and rapeseed oils in cooking is the first step to increasing your intake of these acids. You can also consider taking a 2000-4000mg daily dose of an evening primrose oil supplement after consulting your doctor.

related links

- 📄 ARTICLE: [Are there any herbal remedies that will help me sleep?](#)
- 📄 ARTICLE: [Green-lipped mussels and your health](#)
- 📄 ARTICLE: [Can vitamin E make me look or feel younger?](#)
- 📄 ARTICLE: [Ginkgo biloba – thanks for the memory](#)
- 📄 ARTICLE: [Fish oils help arthritis](#)

your b
Ma
Pe

J
iVilla
date o
for

This iVillage Health service area is designed for educational purposes only. You should not rely on this information as a substitute for medical attention, diagnosis or hands-on treatment. If you are concerned about your health or that of a child, please consult your health provider immediately and do not wait for a response from our professionals. For the full Disclaimer, [click here](#).



Send this page to a friend

Friend's
Email:

Your
Email:

CHANNELS

Computers & Internet
Diet & Fitness
Food
Games
Health & Beauty
Horoscopes & Astrology
Money

Motoring
News & Politics
Pregnancy & Baby
Relationships
Shopping
Work & Career

COMMUNITY

Join free
Message Boards
Member Centre
Resource Centre
Help

SERVICES

Auctions
Competitions
Email
eCards
Newsletters
Phonebook
Quizzes
Site Map
Weather

ABOUT IVILLAGE

Contact Us
Corporate Profile
iVillage.com
Jobs
Mission Statement
Press Releases
Privacy Policy
Terms of Service

iVillage.co.uk	Search	<input type="text"/>	<input type="button" value="Go!"/>	<input type="button" value="Computers Diet & Fitness Food Health Horoscopes iVillage UK Money News Pregnancy Relationships Work & Career the Internet"/>

© iVillage UK Limited 2000-2002. All rights reserved. Terms of Service.

© iVillage inc. 1995-2002. All rights reserved.

Evening Primrose Oil

Also indexed as: Black Currant Seed Oil, EPO, *Oenothera biennis*, Starflower

What does it do? Evening primrose oil (EPO), black currant seed oil, and borage oil contain gamma linolenic acid (GLA), a fatty acid that the body converts to a hormone-like substance called prostaglandin E1 (PGE1). PGE1 has anti-inflammatory properties and may also act as a blood thinner and blood vessel dilator.

The anti-inflammatory properties of EPO have been studied in double-blind research with people suffering from rheumatoid arthritis. Some, but not all, studies have reported that EPO supplementation provides significant benefit to these people.¹




Photo copyright Steven Foster

GLA, the primary active ingredient in EPO, has anticancer activity in test tube studies² and in some,³ but not all,⁴ animal studies. Injecting GLA into tumors has caused regression of cancer in people in preliminary research.⁵ Very preliminary evidence in people with cancer suggested "marked subjective improvement,"⁶ though not all studies find GLA helpful.⁷

EPO has been reported to lower cholesterol levels in people in some,⁸ but not all,⁹ research.

Linoleic acid, a common fatty acid found in nuts and seeds and most vegetable oils (including EPO), should theoretically convert to PGE1. But many things can interfere with this conversion, including disease, the aging process, saturated fat, hydrogenated oils, blood sugar problems, and inadequate vitamin C, magnesium, zinc, and B vitamins. Supplements that provide GLA circumvent these conversion problems, leading to more predictable formation of PGE1.¹⁰

Where is it found? EPO is found primarily in supplements. The active ingredient, GLA, can also be found in black current seed oil and borage oil supplements.

Evening primrose oil has been used in connection with the following conditions (refer to the individual health concern for complete information):

Ranking	Health Concerns
Primary	<u>Diabetes</u> <u>Eczema</u>
Secondary	<u>Alcohol withdrawal</u> <u>Fibrocystic breast disease</u> <u>Osteoporosis</u> (in combination with fish oil) <u>Phenylketonuria</u> (if PUFA deficient) <u>Premenstrual tension</u> (PMT) <u>Rheumatoid arthritis</u>
Other	<u>Atherosclerosis</u> <u>Attention deficit disorder</u> <u>Intermittent claudication</u> <u>Irritable bowel syndrome</u> (IBS) <u>Multiple sclerosis</u> <u>Raynaud's disease</u> Scleroderma Sjogren's syndrome <u>Tardive dyskinesia</u>

Who is likely to be deficient? Those with premenstrual tension,¹¹ diabetes,¹² scleroderma,¹³ Sjogren's syndrome,¹⁴ tardive dyskinesia,¹⁵ eczema,¹⁶ and other skin conditions¹⁷ can have a metabolic block that interferes with the body's ability to make GLA. In preliminary research, supplementation with EPO has helped people with these conditions.^{18 19 20 21 22}

Though preliminary, double-blind evidence suggests that alcoholics may be deficient in GLA and that alcohol withdrawal may be facilitated with EPO supplementation.²³ Many people in Western societies may be at least partially GLA-deficient as a result of aging, glucose intolerance, dietary fat intake, and other problems. Individuals with deficiencies benefit from supplemental GLA intake from EPO, black currant seed oil, or borage oil.

How much is usually taken? Although many people may have inadequate levels of GLA, the optimal intake for this nutrient remains unknown. Researchers often use 3,000–6,000 mg of EPO per day, which provides approximately 270–540 mg of GLA.

Are there any side effects or interactions? EPO has been reported to exacerbate symptoms of temporal lobe epilepsy, which can sometimes be mistaken for schizophrenia.^{24 25}

Other nutrients are needed by the body, along with EPO, to make PGE1. Consequently, some experts suggest that magnesium, zinc, vitamin C, niacin, and vitamin B6 should be taken along with EPO.

At the time of this writing, no evidence of drug interactions with evening primrose oil was found in the medical literature.

Within *Healthnotes Online*, information about the effects of a particular supplement or herb on a particular condition has been qualified in terms of the methodology or source of supporting data (for example: clinical, double blind, meta-analysis, or traditional use). For the convenience of the reader, the information in the table listing the supplements for particular conditions is also categorized. The criteria for the categorizations are: "Primary" indicates there are reliable and relatively consistent scientific data showing a health benefit. "Secondary" indicates there are conflicting, insufficient, or only preliminary studies suggesting a health benefit or that the health benefit is minimal. "Other" indicates that an herb is primarily supported by traditional use or that the herb or supplement has little scientific support and/or minimal proven health benefit.

References:

1. Joe LA, Hart LL. Evening primrose oil in rheumatoid arthritis. *Ann Pharmacother* 1993;27:1475-7 [review].
2. Dippneaar N, Booyens J, Fabbri D, Katzeff IE. The reversibility of cancer: evidence that malignancy in melanoma cells is gamma linolenic acid deficiency-dependent. *SAfr Med J* 1982;62:505-9.
3. Pritchard GA, Mansel RE. The effects of essential fatty acids on the growth of breast cancer and melanoma. In *Omega-6 Essential Fatty Acids: Pathophysiology and Roles in Clinical Medicine*, ed. DF Horrobin. New York: Alan R Liss, 1990, 379-90.
4. Lee JH, Sugano M. Effects of linoleic and gamma-linolenic acid on 7,12-dimethylbenz(a)anthracene-induced rat mammary tumors. *Nutr Rep Internat* 1986;34:1041.
5. Naidu MRC, Das UN, Kshan A. Intratumoral gamma-linolenic acid therapy of human gliomas. *Prostaglandins Leukotrienes Essential Fatty Acids* 1992;45:181-4.
6. Van der Merwe CF, Booyens J. Oral gamma-linolenic acid in 21 patients with untreatable malignancy. An ongoing pilot open clinical trial. *Br J Clin Pract* 1987;41:907-15.
7. McIlmurray MB, Turkie W. Controlled trial of gamma linolenic in Duke's C colorectal cancer. *Br Med J* 1987;294:1260.
8. Ishikawa T, Fujiyama Y, Igarashi O, et al. Effects of gammalinolenic acid on plasma lipoproteins and apolipoproteins. *Atherosclerosis* 1989;75:95-104.
9. Bobberg M, Vessby B, Selinus I. Effects of dietary supplementation with n-6 and n-3 long-chain polyunsaturated fatty acids on serum lipoproteins and platelet function in hypertriglyceridaemic patients. *Acta Med Scand* 1986;220:153-60.
10. Horrobin DF. The importance of gamma-linolenic acid and prostaglandin E1 in human nutrition and medicine. *J Holistic Med* 1981;3:118-39.
11. Horrobin DF, Manku M, Brush M, et al. Abnormalities in plasma essential fatty acid levels in women with pre-menstrual syndrome and with non-malignant breast disease. *J Nutr Med* 1991;2:259-64.
12. Keen H, Payan J, Allawi J, et al. Treatment of diabetic neuropathy with gamma-linolenic acid. *Diabetes Care* 1993;16:8-15 [reviews].
13. Horrobin DF. Essential fatty acid metabolism in diseases of connective tissue with special reference to scleroderma and to Sjogren's syndrome. *Med Hypoth* 1984;14:233-47.
14. Horrobin DF, Campbell A. Sjogren's syndrome and the sicca syndrome: the role of prostaglandin E1 deficiency. Treatment with essential fatty acids and vitamin C. *Med Hypoth* 1980;6:225-32.
15. Vaddadi KS, Gilleard CJ. Essential fatty acids, tardive dyskinesia, and schizophrenia. In *Omega-6 Essential Fatty Acids: Pathophysiology and Roles in Clinical Medicine*, ed. DF Horrobin. New York: Alan R Liss, 1990, 333-43.
16. Manku MS, Horrobin DF, Morse NL, et al. Essential fatty acids in the plasma phospholipids of patients with atopic eczema. *Br J Derm* 1984;110:643.
17. Horrobin DF. Essential fatty acids in clinical dermatology. *J Am Acad Dermatol* 1989;20:1045-53.
18. Mansel RE, Pye JK, Hughes LE. Effects of essential fatty acids on cyclical mastalgia and noncyclical breast disorders. In *Omega-6 Essential Fatty Acids: Pathophysiology and Roles in Clinical Medicine*, ed. DF Horrobin. New York: Alan R Liss, 1990, 557-66.
19. Keen H, Payan J, Allawi J, et al. Treatment of diabetic neuropathy with gamma-linolenic acid. *Diabetes Care* 1993;16:8-15.
20. Horrobin DF. Essential fatty acid metabolism in diseases of connective tissue with special reference to scleroderma and to Sjogren's syndrome. *Med Hypoth* 1984;14:233-47.
21. Vaddadi KS, Gilleard CJ. Essential fatty acids, tardive dyskinesia, and schizophrenia. In *Omega-6 Essential Fatty Acids: Pathophysiology and Roles in Clinical Medicine*, ed. DF Horrobin. New York: Alan R Liss, 1990, 333-43.
22. Schalin-Karrila M, Mattila L, Jansen CT, et al. Evening primrose oil in the treatment of atopic eczema: effect on clinical status, plasma phospholipid fatty acids and circulating blood prostaglandins. *Brit J Dermatol* 1987;117:11-9.
23. Glen AIM, Glen EMT, MacDonnell LEF, et al. Essential fatty acids in the management of withdrawal symptoms and tissue damage in alcoholics, presented at the 2nd International Congress on Essential Fatty Acids, Prostaglandins and Leukotrienes, London, Zoological Society. March 24-7, 1985, abstract 53.
24. Vaddadi KS. The use of gamma-linolenic acid and linoleic acid to differentiate between temporal lobe epilepsy and schizophrenia. *Prostaglandins Med* 1981;6:375-9.
25. Holman CP, Bell AFJ. A trial of evening primrose oil in the treatment of chronic schizophrenia. *J Orthomol Psychiatr* 1983;12:302-4.

healthnotes

Copyright © 2000 Healthnotes, Inc. All rights reserved. www.healthnotes.com

The information presented in *Healthnotes Online* is for informational purposes only. It is based on scientific studies (human, animal, or *in vitro*), clinical experience, or traditional usage as cited in each article. The results reported may not necessarily occur in all individuals. For many of the conditions discussed, treatment with prescription or over-the-counter medication is also available. Consult your doctor, practitioner, and/or pharmacist for any health problem and before using any supplements or before making any changes in prescribed medications. Information expires January 2002.

This is **G o o g l e**'s cache of <http://www.mineralconnection.com/evenprim.htm>.

G o o g l e's cache is the snapshot that we took of the page as we crawled the web.

The page may have changed since that time. Click here for the current page without highlighting.

To link to or bookmark this page, use the following url:


<http://www.google.com/search?q=cache:uK48zjdg4McC:www.mineralconnection.com/evenprim.htm+evening+primrose+oil+omega&hl=en&ie=UTF-8>

Google is not affiliated with the authors of this page nor responsible for its content.

These search terms have been highlighted: **evening primrose oil omega**



Evening Primrose Oil



Country® Life

Evening Primrose Oil

Highest Natural Source of GLA

500 mg

Gamma Linolenic Acid (GLA) and Linoleic Acid are **Omega-6** fatty acids. they are responsible for the production of the PGE-1 family of prostaglandins, the hormone substances which help maintain the balance of many metabolic processes.

Essential fatty acids cannot be manufactured by the body and must be derived from food sources.

Country Life's **Evening Primrose Oil** is imported from Europe and is the finest quality available.

Suggested Use: Adults take 2 to 4 softgel capsules daily or more as directed by a health care professional.

Each Softgel Capsule Contains:	
Evening Primrose Oil	500 mg.
Providing 50 mg. of GLA (gamma-linolenic acid) and 350 mg. of Cis-Linoleic Acid	


No Yeast, Corn, Wheat, Soy, Milk, Salt, Sugar, Starch, Preservatives or Artificial Color.

Guaranteed for freshness, purity and specified content.

Click the button with a picture of a shopping cart and the words: "ADD ADD" to order. [Click here](#) for shipping charges and alternative ways to order. [Click here](#) for a list of items refused by Customs.



1579404001 ~ Evening Primrose Oil 500 mg ~ 60 Capsules ~ \$11.10



Health From The Sun

Evening Primrose Oil

**Hexane-Free
500 mg. each**

Health From The SunTM Evening Primrose Oil is one of the best sources of the nutritionally important omega-6 fatty acids, Gamma-Linolenic Acid [GLA] and Cis-Linoleic Acid. The oil is extracted from the highest quality Evening Primrose seeds using the cold expeller-press method, without the chemical solvent hexane. It is a BIO-EFA® food oil, ensuring the most Bioavailable Essential Fatty Acids. Independently tested to guarantee purity and truth-in-labeling.

Health From The Sun'sTM involvement with evening primrose oil goes back to its beginning as a product in the industry. We were the first company to introduce Evening Primrose Oil products to the United States!

Evening Primrose Oil, more commonly referred to as EPO, is by far the most popular source of Omega-6 EFAs, rich in both Linoleic Acid (LA) and Gamma Linolenic Acid (GLA). More research has been done on EPO than any other nutritional oil. This has contributed to its popularity, particularly in relation to PMS, cardiovascular disease, inflammation, and skin disorders.*




Each Capsule Contains:	
500 mg of 100% pure Evening Primrose Oil, providing:	
Cis-Linoleic Acid (omega-6)	370 mg.
Gamma-Linolenic Acid [GLA] (omega-6)	50 mg.
Oleic Acid (omega-9)	35 mg.

***This statement has not been evaluated by the Food & Drug Administration. This product is not intended to diagnose, treat, cure or prevent any disease. Health From The Sun™, "The Leader in Essential Fatty Acid Nutrition"**

Usage: Three to six capsules daily.

No sugar, starch, artificial preservatives or colors. Keep in a cool dry place. Do not use if safety seal is broken or missing.

Click the button with a picture of a shopping cart and the words: "ADD ADD" to order. [Click here](#) for shipping charges and alternative ways to order. [Click here](#)

1004301073 ~ Evening Primrose Oil 500 mg ~ 50 Capsules ~ \$10.95  <small>10 items refused by Customs.</small>	1004301074 ~ Evening Primrose Oil 500 mg ~ 100 Capsules ~ \$17.95  <small>10 items refused by Customs.</small>	1004301076 ~ Evening Primrose Oil 500 mg ~ 250 Capsules ~ \$33.95  <small>10 items refused by Customs.</small>
--	---	---



Health From The Sun

Evening Primrose Deluxe

Hexane-Free
1,300 mg. each

Promotes Healthy Skin and Hormonal Balance®

Health From The Sun's™ involvement with evening primrose oil goes back to its beginning as a product in the industry. We were the first company to introduce Evening Primrose Oil products to the United States!

Evening Primrose Oil, more commonly referred to as EPO, is by far the most popular source of Omega-6 EFAs, rich in both Linoleic Acid (LA) and Gamma Linolenic Acid (GLA). More research has been done on EPO than any other nutritional oil. This has contributed to its

popularity, particularly in relation to PMS, cardiovascular disease, inflammation, and skin disorders.*

Supplement Facts

Serving Size: 1 Capsule

	Amount Per Serving	% Daily Value
Calories	10	
Calories From Fat	10	
Total Fat	1.5 grams	2%*
Gamma-Linolenic Acid (GLA) (Omega-6)	130 mg	+
Linoleic Acid [GLA] (omega-6)	950 mg	+
Oleic Acid (omega-9)	80 mg.	+
* Percent Daily Values are based on a 2000 calorie diet		
+ Daily Value Not Established		
Ingredients: Evening Primrose Seed Oil, Gelatin, Glycerin and Water		

*This statement has not been evaluated by the Food & Drug Administration. This product is not intended to diagnose, treat, cure or prevent any disease. Health From The Sun™, "The Leader in Essential Fatty Acid Nutrition"

Two capsules of Health From The Sun™ Evening Primrose Deluxe supply more GLA than five standard 500 mg Evening Primrose Oil capsules.


Directions: Two or Three capsules daily.

No sugar, starch, artificial preservatives or colors. Keep in a cool dry place. Do not use if safety seal is broken or missing.

Click the button with a picture of a shopping cart and the words: "ADD ADD" to order. [Click here](#) for shipping charges and alternative ways to order. [Click here](#)

1004301078 ~ Evening
Primrose Oil 1,300 mg
~ 50 Capsules ~ \$17.99

Looking For Som Maybe You'll Fi

<p><u>EFAs and Herbs: A Team Defense Against Disease --</u> The latest word indicates that EFAs and herbs, when combined, create a doubly potent defense team greater than the sum of their parts. Learn why two experts believe blending EFAs and herbs can significantly impact the health benefits of both.</p>	<p>supp</p> <p>Depa</p>
<p> <u>Organic Flax Seed ~</u> Another "Essential" Fatty Acid</p>	<p>Depa</p>
<p><u>Books on The Health Benefits of Flax, Evening Primrose Oil and other Essential Fatty Acids</u></p>	
<p><u>The Women's Department</u></p>	<p>Depa</p>



Please note: This flyer is for information purposes only because it has not been evaluated by the Food and Drug Administration. It is not intended to diagnose or prescribe and these products are not intended to treat, cure prescribe or prevent any disease. If you have a medical condition, please see your health professional. Thank you.

If you don't see what you want, just ask. We carry a lot more than we can possibly list here on the web.


We Cheerfully accept:



along with checks and money orders.

30 Day Money Back Guarantee*

*** Does Not Include Postage**



Here's How to Contact Us:

The Mineral Connection 208 N. Main St. Taylor, TX 76574	Local Phone 512/352-1001 Austin Line 512/365-2300 Toll Free in U.S. 800/742-2560 Fax 512/352-1002	<u>E- Mail</u>
--	--	-----------------------

© Copyright 1998-2002 Glynda Chambliss & The Mineral Connection

Designed 12/06/97

Setup on 03/11/98 ~ Revised 01/07/02

BEV EPO is a rich source of Omega-6 fatty acids. These are considered essential fatty acids (EFA) due to their requirement for proper human growth and development. EPO is one of the richest sources of the Omega-6 fatty acid GLA (Gamma Linolenic Acid). BEV EPO supplies essential nutrition to help support body recomposition, immune system effectiveness, healthy skin, cardiovascular protection, and health and well being.

A QUICK HISTORY OF EPO

The evening primrose is a very unique plant. It reaches a height of eight feet tall and its flowers open only in the evening, fluorescing under the light of the moon. Its healing properties allow it to help strengthen the heart and the circulatory system. The history of the use of evening primrose is not as extensive as many herbal remedies. The Coahuilla Indians gathered leaves and caterpillars from it for food. Early English settlers of the Americas took evening primrose back to England where it was cultivated for its nut-flavored root. It is modern scientific research which has brought the health benefits of this plant to light. It has been discovered that an oil derived from the seeds of this plant is helpful in a wide range of health problems, including obesity, premenstrual syndrome, arthritis, multiple sclerosis, brittle nails, eczema, heart disease, high blood pressure, alcoholism and schizophrenia.

GAMMA LINOLENIC ACID

The key to effectiveness of evening primrose oil is the presence of gamma linolenic acid (GLA). GLA is an essential fatty acid. Essential fatty acids, like vitamins and some amino acids, are substances that the body needs to function properly but that it cannot produce. This means that GLA must come from dietary sources. But what is the function of GLA? GLA is the parent compound to omega-3 and omega-6 fatty acids, but it is even more important in the production of substances known as prostaglandins. Our bodies take linoleic acid (abundant in vegetables) and change it into GLA, then into dihomogamma-linoleic acid to arachadonic acid. The arachadonic acid is then used to produce prostaglandins. Prostaglandins are hormone-like substances that regulate blood pressure and clotting.

WHY DO I NEED A SUPPLEMENT OF GLA?

By now you may be wondering, "if I eat vegetables that supply linoleic acid, why do I need a supplement of GLA?" Our diet, high in saturated fats and processed vegetable oils, inhibits the body's ability to change linoleic acid to GLA. This means that by supplementing the diet with GLA, the body can skip a step and more efficiently produce the prostaglandins. As it turns out, evening primrose is the only significant dietary source of GLA besides black currant oil and breast milk. That is why evening primrose oil can be so important to relieving the conditions already mentioned. One of the overall effects of evening primrose oil is that it works as an anticoagulant to lower blood pressure and strengthen the heart.



PRICE: \$18.00

ORDER NOW!
CALL TOLL FREE
For U.S. Orders Call:
1.800.888.3364

OR
FAX YOUR ORDER
Click Here For
Printable Fax Form
1.361.654.4050

E-Mail

RETAIL ORDERS OF
\$125.00 OR MORE IN
PRODUCT
RECEIVE FREE
SHIPPING*
RETAIL ORDERS
UNDER \$125.00
ADD \$5.00*
***SHIPPING RATES**
FOR THE 48
CONTIGUOUS
STATES ONLY

BEV EPO EVENING PRIMROSE OIL

DIRECTIONS: The generally recommended dosage of BEV EPO is one to three softgel capsules taken daily with meals. One (1) softgel capsule of BEV EPO provides 10 calories from 1000 mg Evening Primrose Oil which typically contains 700 mg Linoleic Acid and 90 mg Gamma Linolenic Acid.

Supplement Facts

Serving Size: 1 Softgel Capsule

Servings Per Container: 90

Amount Per Serving:

Calories: 10

Calories from fat: 10

% Daily Value*

Total Fat	1 g	2%
Polyunsaturated Fat	1 g	**
Evening Primrose Oil	1000 mg	**
Linoleic Acid	700 mg	**
Gamma Linolenic Acid	90 mg	**

*Percent Daily Values are based on a 2000 calorie diet

**Daily Value Not Established

Other Ingredients: Gelatin and Glycerin

Free Shipping is not
included with any
promotions.

WE ACCEPT VISA,
MASTERCARD,
DISCOVER and
NOVUS.

Our No Nonsense

Guarantee: If you are not
totally satisfied with our
products, simply return
the unused portion and we
will cheerfully replace
your order or refund your
money, whichever you
prefer.

[TO ORDER](#)

[PRICE LIST](#)

[HOME PAGE](#)

info@beverlyintl.com



Flaxseed – METABOLISM OF OMEGA-3 AND OMEGA-6 FATTY ACIDS

Fatty Acid Profile of Flaxseed

Flaxseed contains a mix of fatty acids. It is high in polyunsaturated fatty acids (73%), moderate in monounsaturated fatty acids (18%), and low in saturated fatty acids (9%). The saturated fat level of flaxseed is similar to that of canola. Flaxseed is a rich plant source of alpha-linolenic acid (ALA), an essential fatty acid in the human diet and the parent fatty acid of the omega-3 family, as shown in Figure 1. ALA is converted to two main long-chain fatty acids, eicosapentaenoic acid (EPA) and docosahexaenoic acid (DHA), in a series of enzymatic reactions (below). ALA has been shown to modulate eicosanoid synthesis,² and its concentration in breast milk exceeds that of DHA, suggesting a particular requirement for ALA by infants.^{3, 4}

Omega-6/Omega-3 Fatty Acid Ratio of Flaxseed

ALA comprises about 57% of the total fatty acids in flaxseed, whereas the omega-6 fatty acids comprise about 16%, giving an omega-6/omega-3 ratio of 0.3:1.¹ Because the typical Western diet is high in linoleic acid and low in omega-3 fatty acids, some experts recommend replacing omega-6 fatty acids with those from the omega-3 family. Consuming flaxseed or foods rich in ALA, such as omega-3 enriched eggs derived from hens fed flaxseed, increases omega-3 fatty acid intake. This improves the dietary omega-6/omega-3 ratio.⁵

Major Families of Unsaturated Fatty Acids

There are four major families of unsaturated fatty acids (see Table 1). Oleic acid is the most prevalent fatty acid in nature and can be synthesized in the body from dietary stearic acid (18:0).⁶ Palmitoleic acid can be synthesized from dietary palmitic acid (16:0). Oleic and palmitoleic acids are not essential in human nutrition because they can be formed from dietary precursors.

Two fatty acids are required in the diets of humans because our bodies cannot manufacture them from dietary precursors: Alpha-linolenic acid, the parent compound of the omega-3 fatty acid family, and linoleic acid, the parent compound of the omega-6 family. Arachidonic acid, a metabolite of linoleic acid, is considered an essential fatty acid (EFA) only when a linoleic acid deficiency exists.⁷

Desaturation and elongation. Alpha-linolenic acid and linoleic acid are converted to their respective metabolites by a series of alternating desaturations and elongations. The desaturations add a double bond by removing hydrogen, while the elongations add two carbon atoms.

TABLE 1

Nomenclature of Major Families of Unsaturated Fatty Acids^a

Parent Compound	Number of Double Bonds	Family Name ^b	Structural Abbreviations ^c
Oleic acid	one	Omega-9 (ω-9)	18:1n-9 or 18:1ω-9
Palmitoleic acid	one	Omega-7 (ω-7)	16:1n-7 or 16:1ω-7
Linoleic acid	two	Omega-6 (ω-6)	18:2n-6 or 18:2ω-6
Alpha-linolenic acid ^d	three	Omega-3 (ω-3)	18:3n-3 or 18:3ω-3

^a Adapted from Vaisey-Genser M. In: *Flaxseed: Health, Nutrition and Functionality*. Winnipeg, MB: Flax Council of Canada, 1994, p. 11.


^b The family name denotes the position of the first double bond as the number of carbon atoms from the methyl end of the fatty acid chain.

^c Number of carbon atoms: number of double bonds (fatty acid family).

^d Also designated α-linolenic acid. Alpha-linolenic acid is distinct from gamma-linolenic acid (γ-linolenic acid (18:3n-6)), which is an intermediate in the omega-6 metabolic pathway and is a major component of evening primrose, borage and black currant oils.

FIGURE 1

Metabolic Pathways of the Omega-3 and Omega-6 Fatty Acids



Only the names of major fatty acids are shown in this figure.



The first step in the metabolism of both EFA families is desaturation, catalysed by delta-6-desaturase. This step is followed by elongation, then desaturation (catalysed by delta-5-desaturase), then elongation and, finally, desaturation (catalysed by delta-4-desaturase). The desaturation steps tend to be slow, while the elongation steps are rapid. Thus, the tissue concentrations of gamma-linolenic acid (GLA) (18:3n-6) and stearidonic acid (18:4n-3) tend to be low, because they are formed slowly by desaturation and then quickly elongated to other metabolites.^{8,9}

Competition between families. Mammals cannot interconvert the omega-3 and omega-6 fatty acids. Furthermore, their metabolism requires the same desaturation enzymes, resulting in competition between the two families. An excess of one family of fatty acids can interfere with the metabolism of the other, reducing their incorporation into tissue lipids and altering their biological effects.^{8,10}

Metabolism of Alpha-Linolenic Acid (ALA)

Dietary ALA appears to have two metabolic fates. It can be desaturated and elongated to its long-chain metabolites or it can undergo β -oxidation. Although studies of ALA β -oxidation in humans have not been conducted, studies in animals suggest that the metabolism of ALA by β -oxidation may contribute significantly to energy production.¹¹ Additional research is needed to clarify the relative importance of these metabolic pathways.

Conversion of ALA to EPA and DHA. About 15% of ALA is converted to EPA and about 5% is converted to DHA in a process that is somewhat slow in humans.^{5,11} Conversion of ALA to its long-chain metabolites is affected by various dietary factors. A diet rich in linoleic acid, for example, has been shown to reduce ALA conversion by as much as 40%,¹⁰ and a high maternal intake of linoleic acid lowers EPA and DHA levels in umbilical plasma, suggesting reduced ALA conversion and availability of omega-3 fatty acids to the developing fetus.¹² Saturated and *trans* fatty acids also interfere with ALA desaturation and elongation,^{13,14} and in rats, ethanol inhalation results in a significant loss of liver DHA, indicating reduced ALA conversion.¹⁵

DHA can be converted back into EPA in a reaction called retroconversion. This is believed to be a minor metabolic pathway in humans.¹⁶

References

1. Canadian Grain Commission. *Flaxseed Export Quality Data*. Winnipeg, MB.
2. Ferretti A and Flanagan VP. *Prostaglandins Leukot Essent Fatty Acids*. 1996;54:451-455.
3. Ratnayake WMN and Chen Z-Y. *Lipids*. 1996;31(Suppl):S279-S282.
4. Budowski P, et al. *World Rev Nutr Diet*. 1994;75:105-108.
5. Nutrition Advisory Panel Meeting: Executive Summary. Winnipeg, MB: Flax Council of Canada, 1995.
6. *Food Fats and Oils*. Washington, DC: Institute of Shortening and Edible Oils, Inc., 1994.
7. Food and Nutrition Board, National Research Council. In: *Recommended Dietary Allowances*, 10th ed. Washington, DC: National Academy Press, 1989, pp. 44-51.
8. Horrobin DF and Manku MS. In: *Omega-6 Essential Fatty Acids*. Horrobin DF, ed. New York, NY: Alan R. Liss, 1990, pp. 21-53.
9. Salem Jr N and Pawlosky RJ. *World Rev Nutr Diet*. 1994;75:114-119.
10. Emken EA. In: *Proceedings from the Scientific Conference on Omega-3 Fatty Acids in Nutrition, Vascular Biology, and Medicine*. Dallas, TX: American Heart Association, 1995, pp. 9-18.
11. Cunnane SC. In: *Flaxseed in Human Nutrition*. Cunnane SC and Thompson LU, eds. Champaign, IL: AOCS Press, 1995, pp. 99-127.
12. Al MDM, et al. *J Am Coll Nutr*. 1996;15:49-55.
13. Ackman RG and Cunnane SC. In: *Advances in Applied Lipid Research*. Padley FB, ed. London: JAI Press Ltd., 1992, pp. 167-215.
14. Houwelingen AC v and Hornstra G. *World Rev Nutr Diet*. 1994;75:175-178.
15. Salem Jr N, et al. *Lipids*. 1996;31(Suppl):S153-S156.
16. Brossard N, et al. *Am J Clin Nutr*. 1996;64:577-586.